

GENERAL INDEX TO VOLUME III

New scientific names of plants and the final members of new combinations are printed in **bold face type**; synonyms and page numbers having reference to figures or plates, in *italic*; and previously published scientific names and all other matter, in ordinary type.

A

Achorion Schönleinii, 268
Acrostalagmus albus, 6
Agrimonia, 309; *americana*, 313; *Eupatoria*, 313, 315, 316, 317, var. *americana*, 317, var. *glabra*, 313, var. *hirsuta*, 313, var. *mollis*, 315, var. *parviflora*, 313; *gryposepala*, 313; *hirsuta*, 313; *microcarpa*, 315; *mollis*, 316; *parviflora*, 317; *parviflora*, 313, 316; *platycarpa*, 315; *polyphylla*, 317; *pubescens*, 315; *pumila*, 315; *rostellata*, 313; *serrifolia*, 317; *striata*, 313; *suaveolens*, 317
Agrimonia, The Missouri, 309
Alternaria sp., cultures of, 279, 281
Amidases in *Lenzites saepiaria*, 489; in other fungi, 489
Apparatus, a note on the adaptability of the Folin micro-Kjeldahl, for plant work, 407
arachnoidea (*Thelephora*), 213
argillaceus (*Hypochnus*), 222
Ascomycetes, 253, 271; temperature relations of the, 51
Aspergillus, 253, 271; *aviarius*, 253, 271; *bronchialis*, 254, 271; *candidus*, 254, 271; *flavescens*, 254; *flavus*, 254, 271; *fontoyonoti*, 255, 271; *fumigatus*, 254, 271; *glauca*, 255, 271; *malignus*, 256, 272; *niger*, nitrogen fixation in, 423, 430; *nigrescens*, 254; *nigricans*, 256, 272; *repens*, 256, 272; *Tokelau*, 256, 272
atratum (*Septobasidium*), 334
atroruber (*Hypochnus*), 230
atroruber (*Zygodesmus*), 230
aurantiaca (*Tomentella*), 241
aurantiacus (*Hypochnus*), 241
avellaneus (*Hypochnus*), 225
Azotobacter chroococcum, nitrogen fixation in, 423, 431; *vinelandii*, nitrogen fixation in, 423, 431

B

Basidiomycetes, temperature relations of the, 53
bicolor (*Hypochnus*), 229

bicolor (*Zygodesmus*), 227
biennis (*Thelephora*), 213
botryoides (*Hypochnus*), 226
botryoides (*Thelephora*), 226
Botryosporium, 22; *pyramidale*, 22
Botrytis cinerea, 4
Burt, E. A. *Pistillaria* (subg. *Pistillina*) *Thaxteri*, *Burt* n. sp., 403; The *Thelephoraceae* of North America, VI, 203; VII, 319
Bush, B. F. The Missouri *agrimoniae*, 309

C

Cabbage yellows, 25; and the relation of temperature to its occurrence, 25; causal organism of, 28; symptoms of, 27
canadensis (*Hypochnus*), 211
canescens (*Septobasidium*), 342
castaneum (*Septobasidium*), 330
Catalase in *Lenzites saepiaria*, 501; in other fungi, 501
Catalogue of the plants of Jasper County, Missouri, 345
Cellulase in *Lenzites saepiaria*, 474; in other fungi, 471
cervinus (*Hypochnus*), 232
chlorochaetes (*Zygodesmus*), 219
Cineraria lyrata, 99
cinerascens (*Hypochnus*), 233
cirratum (*Septobasidium*), 334
Claudopus subnidulans, 195
Coniophora, 204; *arida*, 222
coriarius (*Hypochnus*), 228
Corticium, 204; *echinosporum*, 237; *ferrugineum*, 208; *fumosum*, 239; *fuscum*, 215; *isabellinum*, 222; *sulphureum*, 239; *thelephoroides*, 235; *umbrinum*, 213
Cotton root rot, disease of, 11
Cryptococcus, 263, 272; *Corsellii*, 263; *degenerans*, 263; *farctinosus*, 264; *Gilchristi*, 264; of *Gotti & Brazzola*, 263; *granulomatogenes*, 264; *hominia*, 264; *linguae-pilosae*, 264; *lithogones*, 261, 265; *niger*, 265; *Tokishigei*, 265; *tumefaciens*, 262
Cyto-hydrolyzing enzymes, 465; effect of, on sound and decayed wood, 484

(513)

D

- Davis, A. R.: A note on the adaptability of the Folin micro-Kjeldahl apparatus for plant work, 407; Duggar, B. M., and, Studies in the physiology of the fungi. I, Nitrogen Fixation, 413
- Diastase in *Lenzites saeplaria*, 464; in other fungi, 464
- Duggar, B. M.: *Rhizoctonia Solani* in relation to the "mopopils" and the "vermehrungspilz," 1; The Texas root rot fungus and its conidial stage, 11; and Davis, A. R., Studies in the physiology of the fungi. I, Nitrogen Fixation, 413

E

- echinosporum* (*Corticium*), 237
- echinosporus* (*Hypochnus*), 237
- elaecodes* (*Hypochnus*), 218
- Emig, W. H. The occurrence in nature of certain yeast-like fungi with reference to their possible pathogenicity in the higher animals, 243
- Emulsin in *Lenzites saeplaria*, 462; in other fungi, 461
- Endomyces, 259; *albicans*, 259, 272; *tropicalis*, 260
- Enzyme activity in *Lenzites saeplaria*, 452
- epigaeus* (*Hypochnus*), 226
- Esterases in *Lenzites saeplaria*, 455; in other fungi, 455

F

- ferruginea* (*Thelephora*), 208
- ferruginea* (*Tomentella*), 208
- ferrugineum* (*Corticium*), 208
- ferrugineus* (*Hypochnus*), 207
- ferruginosa* (*Tomentellina*), 212
- ferruginosus* (*Hypochnus*), 212
- fibrillosus* (*Hypochnus*), 238
- flavovirens* (*Tomentella*), 220
- floridana* (*Thelephora*), 214
- Folin and Farmer, modification of Kjeldahl method of nitrogen determination by, 408; apparatus used in, 407, 418
- Förster method of determining nitrogen, 428
- frustulosa* (*Hymenochaete*), 337
- frustulosum* (*Septobasidium*), 337
- fulgineus* (*Hypochnus*), 232
- fulvo-cinctus* (*Hypochnus*), 228
- fumigatum* (*Septobasidium*), 340
- fumosum* (*Corticium*), 239
- fumosus* (*Hypochnus*), 239

Fungi: gill, New or interesting species of, from Missouri, 195; Studies in the physiology of the, I, 413, II, 439; yeast-like, The occurrence in nature of certain, with reference to their possible pathogenicity in the higher animals, 243

- Fungi Imperfecti, 263, 272; temperature relations of the, 52
- Fusarium, 25; *conglutinans*, 28, field observations of, 56, host relations of, 43, inoculation experiments with, 35, temperature relations of, 32, virulence of cultures of, 39; *orthoceras*, 30
- fusca* (*Odontia*), 239
- fusca* (*Thelephora*), 215
- fusca* (*Tomentella*), 215
- fuscata* (*Hypochnopsis*), 213
- fuscatus* (*Hypochnus*), 213
- fuscum* (*Corticium*), 215
- fuscus* (*Hypochnus*), 215

G

- Gill fungi, New or interesting species of, from Missouri, 195
- Gilman, J. C. Cabbage yellows and the relation of temperature to its occurrence, 25
- Glenspora*, 270, 272; *Graphii*, 270, 272
- Glomerella Gossypii*, nitrogen fixation in, 424, 431
- Grandinia*, 204; *coriaria*, 228; *tabacina*, 218
- granosa* (*Thelephora*), 227
- granosus* (*Hypochnus*), 227
- granulosus* (*Hypochnus*), 218
- granulosus* (*Zygodesmus*), 218
- Greenman, J. M. A new *Senecio* from Jamaica, 201; Monograph of the North and Central American Species of the genus *Senecio*—Part II, 85
- Gymnoascaceae, 267

H

- Hemicellulase in *Lenzites saeplaria*, 479
- Hippuricase in *Lenzites saeplaria*, 492; in other fungi, 491
- hyalina* (*Pistillina*), 405
- Hymenochaete frustulosa*, 337
- Hymenomycete, the smallest known, 403
- Hypochnus*, 203; *argillaceus*, 222; *atroruber*, 230; *aurantiacus*, 241; *avellaneus*, 225; *bicolor*, 229; *botryoides*, 226; *canadensis*, 211; *cervinus*, 232; *cinerascens*, 233; *coriarius*, 228; *echinosporus*, 237; *elaecodes*, 218; *epigaeus*, 226; *ferrugineus*, 207; *ferruginosus*, 212; *fibrillosus*, 238; *fulgineus*, 232; *fulvo-cinctus*, 228; *fumosus*, 239;

fuscatus, 213; *fuscus*, 215; *granosus*, 227; *granulosus*, 218; *isabellinus*, 222; *leprosus*, 223; *obscuratus*, 216; *olivascens*, 220; *pannosus*, 223; *peniophoroides*, 234; *pilosus*, 221; *rubiginosus*, 209; *sitnensis*, 213; *sparsus*, 225; *spiniferus*, 218; *spongiosus*, 216; *subferrugineus*, 210; *subvinosus*, 231; *tabacinus*, 218; *thelephoroides*, 235; *tristis*, 213; *umbrinus*, 213; *zygodesmoides*, 236
Hypochnopsis fuscata, 213

I

Inulinase in *Lenzites saepiararia*, 488; in other plants, 488
 Invertase in *Lenzites saepiararia*, 459
isabellinum (*Corticium*), 222
isabellinus (*Hypochnus*), 222

J

jamaicaense (*Septobasidium*), 333
 Jasper County, Missouri, description of, 349; plants of, 359
Jola, 319

K

Kjeldahl method of determining nitrogen, the Folin modification of the original, 407

L

Lactase in *Lenzites saepiararia*, 458
Langloisii (*Septobasidium*), 335
Lenzites saepiararia Fries, 439; growth of, on resin agar plates, 512; pine blocks decayed by, 510; sporophores of, 510; with special reference to enzyme activity, 439
Leprieurii (*Septobasidium*), 332
Lepidophyton, 257
leprosus (*Hypochnus*), 223
 Ligninase in *Lenzites saepiararia*, 469; in other fungi, 466
llacinum (*Septobasidium*), 343

M

Macro-Kjeldahl method of determining nitrogen, 410, 426
 Macrosporium commune, nitrogen fixation in, 423, 430
 Maltase in *Lenzites saepiararia*, 457
Montiopsis Aderholdii, 2
 "Mopopilz" and the "vermehrungspilz,"
Rhizoctonia Solani in relation to the,
 1

Mortierella, 253, 271
Mucor, 250, 270; *cornealis*, 250, 270; *corymbifer*, 250, 270; *Mucedo*, 251, 271; *parasiticus*, 251, 271; *pusillus*, 252, 271; *racemosus*, 252, 271; *ramosus*, 252, 271; *Regnieri*, 251, 271; *septatus*, 252, 271; *Truchisi*, 251, 271

N

Nitrogen fixation, 413; methods of determining amount of, 426; organisms used in experiments in, 423; results of experiments in, 429
 Nuclease in *Lenzites saepiararia*, 493; in other fungi, 493

O

obscuratus (*Hypochnus*), 216
Odontia, 204; *fusca*, 239; *porriginis*, 268
olivacea β *botryoides* (*Thelephora*), 226
olivascens (*Hypochnus*), 220
olivascens (*Zygodesmus*), 221
Oospora, 266, 272; *buccalis*, 267; *lingualis*, 265; *porriginis*, 268; *pulmonalis*, 266; sp., culture of, 281; *tonsurans*, 268; of *Trichophyton*, 267
Otomyces purpureus, 256
 Overholts, L. O. New or interesting species of gill fungi from Missouri, 195
 Oxidases in *Lenzites saepiararia*, 501; in other fungi, 501
Ozonium, 11; *auricomum*, 11; *omnivorum*, 11

P

Palmer, E. J. Catalogue of the plants of Jasper County, Missouri, 345
Panaeolus reticulatus, 195, 200; *rufus*, 196, 200; *variabilis*, 197, 200
pannosus (*Hypochnus*), 223
pannosus (*Zygodesmus*), 223
Patouillardii (*Septobasidium*), 332
 Pectase in *Lenzites saepiararia*, 483; in other fungi, 481
 Pectinase in *Lenzites saepiararia*, 483; in other fungi, 481
pedicellata (*Thelephora*), 323, 324, 327
pedicellatum (*Septobasidium*), 323
Penicillium digitatum, nitrogen fixation in, 423, 431; *expansum*, nitrogen fixation in, 423, 431
Peniophora, 204
peniophoroides (*Hypochnus*) 234
Phlebia vaga, 239
 Phoma Betae, nitrogen fixation in, 423, 431

Phycomycetes, 250, 270; temperature relations of the, 49
 Phymatotrichum, 22; *omnivorum*, 22, host plants of, 12, mycelium of, 18; pyramidale, 22
pilosus (Hypochnus), 221
Pinus echinata, 453; *palustris*, 453; *Taeda*, 453
Pistillaria Thaxteri, 406
Pistillina hyalina, 405
plumbea (Sebacina), 241
plumbescens (Sebacina), 241
Pluteus cervinus var. *caespitosus*, 199, 200
 Protease in *Lenzites saepiaria*, 497; in other fungi, 495
pseudopedicellatum (Septobasidium), 327
punicea (Thelephora), 229

R

Raffinase in *Lenzites saepiaria*, 460; in other fungi, 461
 Rennet in *Lenzites saepiaria*, 500; in other plants, 500
 retiforme (Septobasidium), 338
 retiformis (Thelephora), 338
 Rhinotrichum, 22
 Rhizoctonia Solani in relation to the "mopopile" and the "vermehrungspilz," 1
Rhizomorcor parasiticus, 251; *septatus*, 252
Rhizopus, 253, 271; *Cohnii*, 253, 271; *niger*, 253, 271; *nigricans*, 252, 271
rubiginosa (Tomentella), 209
rubiginosus (Hypochnus), 209
rubiginosus (Zygodesmus), 209

S

Saccharomyces, 257; *anginae*, 257, 272; *cerevisiae*, culture of, 284; *granulatus*, 258, 272
Saccharomycopsis, 258, 272; *guttulatus*, 258, 272
 Schizomycetes, temperature relations of the, 49
 Schweinitzii (Septobasidium), 324
Sclerotinia, 5
Sebacina plumbea, 241; *plumbescens*, 241
 Senecio, A new, from Jamaica, 201; Monograph of the North and Central American Species of the genus—Part II, 85
 Senecio, 85; *acutidens*, 178; *Adamsi*, 172; *anacletus*, 183, 192; *aquariensis*, 144; *aueticus*, 102; *aureus*, 119, var. *gracilis*, 128, var. *semi-*

cordatus, 129; *aureus*, 91, 129, var. *angustifolius*, 155, var. *Balsamitae*, 130, 149, 155, 159, $\geq \times$ *Balsamitae*, 129, var. *borealis*, 91, 96, 170, 174, 179, var. *compactus*, 179, var. *croceus*, 142, 144, 173, var. *discoidea*, 113, var. *discoideus*, 91, 96, var. *lancoelatus*, 130, var. *obovatus*, 108, 174, var. *pauciflorus*, 91, var. *subnudus*, 103; *balsamitae*, 148; *Balsamitae*, 155, 159, 167, var. *Crawfordii*, 139, var. *praelongus*, 159; *Burkei*, 136; *camporum*, 149; *Cardamine*, 116; *compactus*, 179; *condensatus*, 179; *conterminus*, 101; *Crawfordii*, 139; *croceus*, 141, var. *Wolffii*, 143; *croceus*, 145; *cyclophyllus*, 117; *cymbalarioides*, 174, var. *borealis*, 177, var. *streptanthifolius*, 177; *cymbalarioides*, 104, 173, 178, var. *diversilobus*, 171; *debilis*, 96; *densus*, 179; *dimorphophyllus*, 144; *discoideus*, 90, 91, 97; *Earlei*, 155; *Elliottii*, 108; *elongatus*, 91, 113; *Farriae*, 146; *fastigiatus*, 119; *fedifolius*, 90; *Fernaldii*, 90; *flavovirens*, 167, var. *thomsonienseis*, 169; *flavovirens*, 96; *flavulus*, 147, 167; *fulgens*, 167; *gaspensis*, 138; *Grayii*, 107; *Hartianus*, 147; *hesperis*, 102; *hesperius*, 102, 190; *heterodoxus*, 144; *Hollickii*, 201; *hyperborealis*, 98; *hyperborealis* var. *columbiensis*, 100; *idahoensis*, 96; *indecorus*, 91; *Jonesii*, 173; *laetiflorus*, 170; *Lemberti*, 91; *Lindheimeri*, 114; *longipetiolatus*, 142; *Lyallii*, 101; *manitobensis*, 179; *Memmingeri*, 155; *Metcalfei*, 179; *microdontus*, 183; *multnomensis*, 169; *mutabilis*, 179; *nephrophyllus*, 97; *Newcombei*, 103; *oblancoelatus*, 179; *obovatus*, 108, var. *divisifolius*, 112, var. *elongatus*, 113, var. *rotundus*, 114, var. *umbratilis*, 115; *obtusatus*, 108; *ovinus*, 100; *Pammelii*, 118; *pauciflorus*, 91, var. *fallax*, 95; *pauperculus*, 159, var. *firmitfolius*, 166; *plattensis*, 148; *platylobus*, 141; *Porteri*, 106; *prionophyllus*, 140; *pseudoreus*, 132, var. *flavulus*, 136; *pseudotomentosus*, 149; *pyrolloides*, 102; *pyrrhocrous*, 142; *quaerens*, 140; *quebradensis*, 117; *remifolius*, 179; *renifolius*, 106; *resedifolius*, 99, var. *columbiensis*, 100; *resedifolius*, 98, 100; *Robbinsii*, 130; *Rosei*, 105, 190; *rotundifolius*, 119; *rotundus*, 114; *rubricaulis*, 173; var. *aphanactis*, 174; *Rydbergii*, 167; *semi-cordatus*, 129; *Smallii*, 155; *Soldanella*, 107; *suavis*, 179; *subcuneatus*, 174; *subnudus*, 103; *Suksdorfii*, 172;

- Swartzii, 201; toluccanus, 186, 194, var. modestus, 188; *Toluccanus* var. *microdontus*, 183; *Tracyi*, 142; tridenticulatus, 179; *tussilaginoidea*, 119; Wardii, 183, 190; Willingii, 154; *Wootonii*, 183
- Septobasidium, 319; atratum, 334; *canescens*, 342; *castaneum*, 330; *cirratum*, 334; *frustulosum*, 337; *fumigatum*, 340; *jamaicaense*, 333; *Langloisii*, 335; *Leprieurii*, 332; *lilacinum*, 343; *Patouillardii*, 332; *pedicellatum*, 323; *pseudopedicellatum*, 327; *retiforme*, 338; *Schweinitzii*, 324; *Spongia*, 339; *sublilacinum*, 331; *tropicale*, 326
- sitnensis* (*Hypochnus*), 213
- sparsus* (*Hypochnus*), 225
- spiniferus* (*Hypochnus*), 218
- Spongia* (*Septobasidium*), 339
- Spongia* (*Thelephora*), 339
- spongiosa* (*Thelephora*), 216
- spongiosus* (*Hypochnus*), 216
- Sporotrichum*, 268, 272; (*Microsporon*)
- Audouini, 266, 268; *Furfur*, 269; *minutissimum*, 269
- Sterigmatocystis*, 257, 272; *nigra*, 257, 272; *pseudonigra*, 257
- Studies in the physiology of the fungi. I, Nitrogen Fixation, 413; II, *Lenzites saeiparia*, with special reference to enzyme activity, 439
- subferrugineus* (*Hypochnus*), 210
- sublilacina* (*Thelephora*), 331
- sublilacinum* (*Septobasidium*), 331
- subvinosus* (*Hypochnus*), 231
- sulphureum* (*Corticium*), 239
- T**
- tabacinus* (*Hypochnus*), 218
- Tannase in *Lenzites saeiparia*, 463; in other fungi, 463
- Temperature, the relation of: to the occurrence of cabbage yellows, 32; to the Ascomycetes, 51; to the Basidiomycetes, 53; to the Fungi Imperfecti, 52; to the Phycomycetes, 49; to the Schizomycetes, 49
- Texas root rot fungus and its conidial stage, The, 11
- Thaxteri* (*Pistillaria*), 406
- Thelephora*, 204; *arachnoidea*, 213; *biennis*, 213; *botryoides*, 226; *ferruginea*, 208; *floridana*, 214; *fusca*, 215; *granosa*, 227; *olivacea* β *T. botryoides*, 226; *pedicellata*, 323, 324, 327; *punicea*, 229; *retiformis*, 338; *Spongia*, 339; *spongiosa*, 216; *sublilacina*, 331; *umbrina*, 213; *vinosa*, 215; *zygodesmoides*, 236
- Thelephoraceae of North America, The, VI, 203; VII, 319
- Toile, maladie de la, 4
- Tomentella flavovirens*, 220
- Tomentella*, 203; *aurantiaca*, 241; *cinerascens*, 233; *ferruginea*, 208; *fusca*, 215; *rubiginosa*, 209; *tristis*, 213
- Tomentellina*, 203; *ferruginosa*, 212
- Torula neoformans*, 261; sp., cultures of, 275, 281
- tristis* (*Hypochnus*), 213
- tristis* (*Tomentella*), 213
- tropicale* (*Septobasidium*), 326
- U**
- umbrina* (*Thelephora*), 213
- umbrinum* (*Corticium*), 213
- umbrinus* (*Hypochnus*), 213
- Urease in *Lenzites saeiparia*, 489; in other fungi, 489
- V**
- vaga* (*Phlebia*), 239
- "Vermehrungspilz" and the "mopopilz," the relation of *Rhizoctonia Solani* to the, 1
- Verticillium Graphii*, 270
- vinosa* (*Thelephora*), 215
- Y**
- Yeast-like fungi, The occurrence in nature of certain, with reference to their possible pathogenicity in the higher animals, 243
- Z**
- Zeller, S. M. Studies in the physiology of the fungi. II, *Lenzites saeiparia* Fries, with special reference to enzyme activity, 439
- Zygodesmus*, 204; *atroruber*, 230; *bicolor*, 227; *chlorochaetes*, 219; *granulosus*, 218; *olivascens*, 221; *pannosus*, 223; *rubiginosus*, 209, 211
- zygodesmoides* (*Hypochnus*), 236
- zygodesmoides* (*Thelephora*), 236